

Gyrokinetic Particle-In-Cell Calculations of Ion Temperature Gradient Driven Turbulence with Parallel Nonlinearity and Strong Flow Corrections*

James C. Kniep, Jean-Noel Leboeuf and Viktor K. Decyk
Department of Physics and Astronomy
University of California at Los Angeles
Los Angeles, CA 90095-1547

Abstract

Nonlinear gyrokinetic calculations have been performed with the three-dimensional, global, toroidal, particle-in-cell, delta-f, UCAN code¹ to study the effects of parallel nonlinearity and strong (externally imposed) flow corrections on ion temperature gradient driven turbulence. These calculations are electrostatic and adiabatic electrons are used. An equilibrium ion temperature gradient only is imposed, along with a generic parabolic q-profile appropriate to a large aspect ratio tokamak. The strong flow corrections derived by T. S. Hahm² have been added to the usual orbit and weight equations used in the UCAN code. The parallel nonlinearity³ has been reactivated as well in the same equations. To examine the effect of strong flow corrections, the calculations have been performed with fluctuations-generated flows or zonal flows arbitrarily set to zero but with externally imposed sheared flows obeying a variety of profiles more or less localized in radius and with various flow magnitudes. The calculations show that the strong flow corrections have a qualitative effect on the saturation level of the fluctuations and on the steady-state heat flux even at the highest flow magnitudes used. The re-activated parallel nonlinearity, with no externally imposed flow but with fluctuations-generated or zonal flows allowed to self-consistently evolve in the calculations, does however make a significant quantitative difference in the saturation level of the fluctuations and in the saturated flux. The presence of the parallel nonlinearity leads to a reduction in both, in apparent agreement with the results of Villard and colleagues³.

¹ R. D. Sydora, V. K. Decyk, and J. M. Dawson, "Fluctuation-induced heat transport results from a large global 3D toroidal particle model", *Plasma Physics and Controlled Fusion* **38**, A281-94 (1996); J.-N. Leboeuf, J. M. Dawson, V. K. Decyk, M. W. Kissick, T. L. Rhodes, and R. D. Sydora, "Effect of externally imposed and self-generated flows on turbulence and magnetohydrodynamic activity in tokamak plasmas", *Phys. Plasmas* **7**, 1795-1801 (2000).

² T. S. Hahm, "Nonlinear gyrokinetic equations for turbulence in core transport barriers", *Phys. Plasmas* **3**, 4658-4664 (1996).

³ L. Villard, S. J. Allfrey, A. Botino, M. Brunetti, G. L. Falchetto, V. Grandgirard, R. Hatzky, J. Nuhrenberg, A. G. Peeters, O. Sauter, S. Sorge, and J. Vaclavik, "Full radius linear and nonlinear gyrokinetic simulations for tokamaks and stellarators: Zonal flows, applied ExB flows, trapped electrons and finite beta", Paper **TH/1-3**, 19th IAEA Fusion Energy Conference, Lyon, France, 14 - 19 October 2002. Submitted to *Nuclear Fusion*.

* Work supported at UCLA by USDOE Grant No. DE-FG03-01ER54617. This work is also part of the SciDAC Plasma Microturbulence Project.